

Dyons near the Planck scale

Larisa Laperashvili ^{1,2} * and C.R. Das ¹ †

¹ *The Institute of Mathematical Sciences, Chennai, India*

² *The Institute of Theoretical and Experimental Physics, Moscow, Russia*

arXiv:hep-ph/0606043v4 29 Nov 2006

* laper@itep.ru, laper@imsc.res.in

† crdas@imsc.res.in

Abstract

In the present letter we suggest a new model of preons-dyons making composite quark-leptons and bosons, described by the supersymmetric string-inspired flipped $E_6 \times \widetilde{E}_6$ gauge group of symmetry. This approach predicts the possible extension of the Standard Model to the Family replicated gauge group model of type $G^{N_{fam}}$, where N_{fam} is the number of families and G is the symmetry group: $G = SMG$, $SU(5)$, $SO(10)$, E_6 , etc. Here E_6 and $\widetilde{E}_6$ are non-dual and dual sectors of theory with hyper-electric g and hyper-magnetic $\tilde{g}$ charges, respectively. Starting with an idea that the most realistic model leading to the unification of all fundamental interactions (including gravity) is the “heterotic” string-derived flipped model, we have assumed that at high energies $\mu > 10^{16}$ GeV there exists the following chain of the flipped models:

$$SU(3)_C \times SU(2)_L \times U(1)_Y \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X \rightarrow$$

$$SU(5) \times U(1)_X \rightarrow SU(5) \times U(1)_{Z1} \times U(1)_{X1} \rightarrow SO(10) \times U(1)_{X1} \rightarrow E_6,$$

ended by the flipped E_6 gauge group of symmetry at the scale $M_{SSG} \sim 10^{18}$ GeV. Suggesting $\mathbf{N} = 1$ supersymmetric $E_6 \times \widetilde{E}_6$ preonic model we have considered preons as dyons confined by hyper-magnetic strings in the region of energies $\mu \lesssim M_{Pl}$. Our model is based on the recent theory of composite non-Abelian flux tubes in SQCD – analog ANO-strings. Considering the breakdown of E_6 and $\widetilde{E}_6$ at the Planck scale into the $SU(6) \times U(1)$ gauge group, we have shown that the six types of k -strings – composite $\mathbf{N} = 1$ supersymmetric non-Abelian flux tubes – are created by the condensation of spreons-dyons near the Planck scale and have six fluxes quantized according to the Z_6 center group of $SU(6)$: $\Phi_n = n\Phi_0$ ($n = \pm 1, \pm 2, \pm 3$). These fluxes give three types of k -strings with tensions $T_k = kT_0$, where $k = 1, 2, 3$, and produce three (and only three) generations of composite quark-leptons and bosons giving a very specific type of “horizontal symmetry”. Thus, the present model predicts $N_{gen} = N_{fam} = 3$. It was shown that our preonic strings are very thin, with radius $R_{str} \sim 10^{-18}$ GeV $^{-1}$, and their tension T_0 is enormously large: $T_0 \sim 10^{38}$ GeV 2 . It was shown that the condensation of spreons near the Planck scale gives the phase transition at some scales M_{crit} and $\widetilde{M}_{crit}$, which correspond to the following breakdowns of E_6 (or $\widetilde{E}_6$) for preons: $E_6 \rightarrow SU(6) \times U(1)$, or $\widetilde{E}_6 \rightarrow \widetilde{SU}(6) \times \widetilde{U}(1)$. We have calculated the critical values of gauge coupling constants: $\alpha^{-1}(M_{crit}) \approx 4.23$ and $\alpha^{-1}(\widetilde{M}_{crit}) \approx 2.13$. It was investigated that in our world we have quark-leptons and gauge bosons A_μ in the region of energies $\mu \lesssim M_{Pl}$, but monopolic “quark-leptons” and dual gauge fields $\widetilde{A}_\mu$ exist in the region $\mu \gtrsim M_{Pl}$.

Contents:

1. Introduction: Superstring theory and the ‘flipped’ E_6 -unification of gauge interactions.
2. Supersymmetric $E_6 \times \widetilde{E}_6$ preonic model of composite quark-leptons and bosons.
3. Preons are dyons confined by hyper-magnetic strings.
4. The breakdown of E_6 and $\widetilde{E}_6$ groups near the Planck scale.
5. Condensation of spreons near the Planck scale.
6. $N = 1$ supersymmetric non-Abelian flux tubes confined preons.
7. Three generations of composite quark-leptons and bosons.
8. Family replicated gauge group $[E_6]^3$ near the Planck scale.

1. Superstring theory is a paramount candidate for the ultimate theory unifying all fundamental interactions including gravity. It was shown in Refs. [1] that superstrings are free of gravitational and Yang-Mills anomalies if a gauge group of symmetry is $SO(32)$, or $E_8 \times E_8$. The “heterotic” superstring theory $E_8 \times E'_8$ was suggested in [1] as a more realistic model for unification. This ten-dimensional Yang-Mills theory can undergo spontaneous compactification for which E_8 group is broken to E_6 in four dimensions, but E'_8 group remains unbroken and gives “hidden sector” of SUGRA.

In the present investigation we develop a new preonic $E_6 \times \widetilde{E}_6$ model of composite quark-leptons and bosons, in which preons are dyons confined by hyper-magnetic strings. Here E_6 and $\widetilde{E}_6$ are non-dual and dual sectors of theory with hyper-electric g and hyper-magnetic $\tilde{g}$ charges, respectively.

Pati was first [2] who suggested to use the strong magnetic force to bind preons-dyons making the composite objects. This idea has an extension in our model [3] (see also the talk [4]), which was constructed in the light of recent investigations of composite non-Abelian flux tubes in SQCD [5–7].

We start with the ‘flipped’ supersymmetric group of symmetry $E_6 \times \widetilde{E}_6$ and show that the dual sector of this theory described by the group $\widetilde{E}_6$ is broken in our world up to the Planck scale $M_{Pl} \approx 1.22 \cdot 10^{19}$ GeV. The breakdown of the dual sector gives a very specific type of the “horizontal symmetry” predicting three generations of the Standard Model.

2. In Ref. [8] we have considered that only ‘flipped’ $SU(5)$ unifies $SU(3)_C$ and $SU(2)_W$ of the Standard Model (SM) at the GUT scale $M_{GUT} \sim 10^{16}$ GeV. An explanation of the discrepancy between the unification scale M_{GUT} and string scale $M_{str} \sim 10^{18}$ GeV was given by the assumption that there exists a chain of extra intermediate symmetries between M_{GUT} and M_{Pl} :

$$\begin{aligned} SU(5) \times U(1)_X &\rightarrow SU(5) \times U(1)_{Z1} \times U(1)_{X1} \rightarrow \\ &SO(10) \times U(1)_{X1} \rightarrow E_6. \end{aligned} \tag{1}$$

We have considered such Higgs boson contents of the $SU(5)$ and $SO(10)$ gauge groups, which give the flipped E_6 final unification at the scale $\sim 10^{18}$ GeV and decreased running of the inversed gauge coupling constant α^{-1} near the Planck scale. Such an example, presented by Fig. 1, suits the purposes of our new model of preons. Here and below we consider the flipped models in which $SU(5)$ contains Higgs bosons h, h^c and H, H^c belonging to $5_h + \bar{5}_h$ and $10_H + \bar{10}_H$ representations of $SU(5)$, respectively, also 24-dimensional adjoint Higgs field A and Higgs bosons belonging to additional higher representations. Correspondingly, the flipped $SO(10)$ (coming at the superGUT scale M_{SG}) contains $10_h + \bar{10}_h$ and $45_H + \bar{45}_H$, 45-dimensional adjoint A and higher representations of Higgs bosons. As it was shown in [8], such Higgs boson contents lead to the flipped E_6 final unification at the supersuperGUT scale $M_{SSG} \sim 10^{18}$ GeV.

Fig. 1 presents an example of running of the inversed gauge coupling constants $\alpha_i^{-1}(\mu)$ (μ is the energy scale) for $i = 1, 2, 3, X, Z, X1, Z1, 5, 10$. It was shown that

at the scale $\mu = M_{GUT}$ the flipped $SU(5)$ undergoes the breakdown to the supersymmetric (MSSM) $SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X$ gauge group of symmetry, which is the supersymmetric extension of the MSSM originated at the seesaw scale $M_{SS} \approx 10^{11}$ GeV, where heavy right-handed neutrinos appear. A singlet Higgs field S provides the following breakdown to the SM (see [9]):

$$SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y. \quad (2)$$

Supersymmetry extends the conventional SM beyond the scale M_{SUSY} . In our example $M_{SUSY} = 10$ TeV.

The final unification E_6 assumes the existence of three 27-plets of E_6 containing three generations of quarks and leptons including the right-handed neutrinos N_i^c (here $i = 1, 2, 3$ is the index of generations). Quarks and leptons of the fundamental 27 representation decompose under $SU(5) \times U(1)_X$ subgroup as follows:

$$27 \rightarrow (10, 1) + (\bar{5}, -3) + (\bar{5}, 2) + (5, -2) + (1, 5) + (1, 0). \quad (3)$$

The first and second quantities in the brackets of Eq. (3) correspond to the $SU(5)$ representation and $U(1)_X$ charge, respectively. We consider charges Q_X and Q_Z in the units $1/\sqrt{40}$ and $\sqrt{3/5}$, respectively, using assignments: $Q_X = X$ and $Q_Z = Z$ [8].

The conventional SM family which contains the doublets of left-handed quarks Q and leptons L , right-handed up and down quarks u^c , d^c , also e^c , is assigned to the $(10, 1) + (\bar{5}, -3) + (1, 5)$ representations of the flipped $SU(5) \times U(1)_X$, along with right-handed neutrino N^c . These representations decompose under

$$SU(5) \times U(1)_X \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X. \quad (4)$$

This decomposition for the E_6 27-plet is given by Dr C.R. Das in his talk [4].

It is necessary to notice that the flipping of our models:

$$d^c \leftrightarrow u^c, \quad N^c \leftrightarrow e^c \quad (5)$$

distinguishes our ‘flipped $SU(5)$ ’ from the standard Georgi-Glashow $SU(5)$.

We have the following 16 spinorial representation of $SO(10)$:

$$F(16) = F(10, 1) + F(\bar{5}, -3) + F(1, 5). \quad (6)$$

Higgs chiral superfields occupy the 10 representation of $SO(10)$:

$$h(10) = h(5, -2) + h^c(\bar{5}, 2). \quad (7)$$

3. Why three generations exist in Nature ? We suggest an explanation considering a new preonic model of composite SM particles. The model starts from the supersymmetric flipped $E_6 \times \widetilde{E}_6$ gauge group of symmetry for preons.

Considering the $\mathbf{N} = 1$ supersymmetric flipped $E_6 \times \widetilde{E}_6$ gauge theory for preons in $4D$ -dimensional space-time, we assume that preons P and antipreons P^c are dyons with charges g and $\tilde{g}$, respectively, resided in the $4D$ hypermultiplets $\mathcal{P} = (P, P^c)$ and $\tilde{\mathcal{P}} = (\tilde{P}, \tilde{P}^c)$. Here “ $\tilde{P}$ ” designates spreons, but not the belonging to $\widetilde{E}_6$.

The dual sector $\widetilde{E}_6$ is broken in our world to some group $\tilde{G}$, and preons and spreons transform under the hyper-electric gauge group E_6 and hyper-magnetic gauge group $\tilde{G}$ as their fundamental representations:

$$P, \tilde{P} \sim (27, N), \quad P^c, \tilde{P}^c \sim (\overline{27}, \bar{N}), \quad (8)$$

where N is the N -plet of $\tilde{G}$ group. We also consider scalar preons and spreons as singlets of E_6 :

$$P_s, \tilde{P}_s \sim (1, N), \quad P_s^c, \tilde{P}_s^c \sim (1, \bar{N}), \quad (9)$$

which are actually necessary for the entire set of composite quark-leptons and bosons. This idea was suggested in Ref. [10].

The hyper-magnetic interaction is assumed to be responsible for the formation of E_6 fermions and bosons at the compositeness scale Λ_s . The main idea of the present investigation is an assumption that preons-dyons are confined by hyper-magnetic supersymmetric non-Abelian flux tubes which are a generalization of the well-known Abelian Abrikosov-Nielsen-Olesen (ANO)-strings for the case of the supersymmetric non-Abelian theory developed in Refs. [5–7]. As a result, in the limit of infinitely narrow flux tubes (strings) we have the following bound states:

i. quark-leptons (fermions belonging to the E_6 fundamental representation):

$$Q^a \sim P^{aA}(y) \left[\mathcal{P} \exp \left(i\tilde{g} \int_x^y \tilde{A}_\mu dx^\mu \right) \right]_A^B (P_s^c)_B(x) \sim 27, \quad (10)$$

$$\bar{Q}_a \sim (P_s^c)^A(y) \left[\mathcal{P} \exp \left(i\tilde{g} \int_x^y \tilde{A}_\mu dx^\mu \right) \right]_A^B P_{aB}(x) \sim \overline{27}, \quad (11)$$

where $a \in 27$ -plet of E_6 , $A, B \in N$ -plet of $\tilde{G}$, and $\tilde{A}_\mu(x)$ are dual hyper-gluons belonging to the adjoint representation of $\tilde{G}$;

ii. “mesons” (hyper-gluons and hyper-Higgses of E_6):

$$\begin{aligned} M_b^a &\sim P^{aA}(y) \left[\mathcal{P} \exp \left(i\tilde{g} \int_x^y \tilde{A}_\mu dx^\mu \right) \right]_A^B (P^c)_{bB}(x) \\ &\sim 1 + 78 + 650 \quad \text{of } E_6, \end{aligned} \quad (12)$$

$$S \sim (P_s)^A(y) \left[\mathcal{P} \exp \left(i\tilde{g} \int_x^y \tilde{A}_\mu dx^\mu \right) \right]_A^B (P_s^c)_B(x) \sim 1, \quad (13)$$

iii. “baryons” of $\tilde{G}$ -triplet (see below):

$$D_1 \sim \epsilon_{ABC} P^{aA'}(z) P^{bB'}(y) P^{cC'}(x) \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^z \tilde{A}_\mu dx^\mu \right) \right]_{A'}^A \times \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^y \tilde{A}_\mu dx^\mu \right) \right]_{B'}^B \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^x \tilde{A}_\mu dx^\mu \right) \right]_{C'}^C, \quad (14)$$

$$D_2 \sim \epsilon_{ABC} P^{aA'}(z) P^{bB'}(y) (P_s)^{C'}(x) \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^z \tilde{A}_\mu dx^\mu \right) \right]_{A'}^A \times \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^y \tilde{A}_\mu dx^\mu \right) \right]_{B'}^B \left[\mathcal{P} \exp \left(i\tilde{g} \int_X^x \tilde{A}_\mu dx^\mu \right) \right]_{C'}^C, \quad (15)$$

and their conjugate particles.

The bound states (10)–(15) are shown in Fig. 3 as unclosed strings (a) and “baryonic” configurations (b). It is easy to generalize Eqs. (10)–(15) for the case of string constructions of superpartners – squark-sleptons, hyper-gluinos and hyper-higgsinos. Closed strings – gravitons – are presented in Fig. 3(c). All these bound states belong to the E_6 representations and they are in fact the $\mathbf{N} = 1$ 4D superfields.

We assume that near the Planck scale preonic E_6 can be broken by Higgses belonging to the 78-dimensional representation of E_6 (see Ref. [3]):

$$E_6 \rightarrow SU(6) \times SU(2) \rightarrow SU(6) \times U(1), \quad (16)$$

where $SU(6) \times U(1)$ is the largest relevant invariance group of the 78.

If $SU(6) \times U(1)$ group of symmetry works near the Planck scale, then we deal just with the theory of non-Abelian flux tubes in $\mathbf{N} = 1$ SQCD, which was developed recently in Refs. [5–7].

Let us consider the condensation of spreons-dyons at the Planck scale. One can combine the Z_6 center of $SU(6)$ with the elements $\exp(i\pi) \in U(1)$ to get topologically stable string solutions possessing both windings, in $\widetilde{SU(6)}$ and $\widetilde{U(1)}$. Now onwards we assume the dual sector of theory described by $\widetilde{SU(6)} \times \widetilde{U(1)}$, which is responsible for hyper-magnetic fluxes. Then, according to the results obtained in Refs. [5–7], we have a nontrivial homotopy group:

$$\pi_1 \left(\frac{SU(6) \times U(1)}{Z_6} \right) \neq 0, \quad (17)$$

and flux lines form topologically non-trivial Z_6 strings.

Besides $SU(6)$ and $U(1)$ gauge bosons, the model contains six scalar fields charged with respect to $U(1)$ and belong to the 6-plet of $SU(6)$. Considering scalar fields of spreons

$$\tilde{P} = \{ \phi^{aA} \}, \quad (18)$$

which have indices a of $SU(6)$ and A of $\widetilde{SU(6)}$ fundamental multiplets, we construct condensation of spreons in vacuum:

$$\tilde{P}_{vac} = \langle \tilde{P}^{aA} \rangle = v \cdot \text{diag}(1, 1, \dots, 1), \quad a, A = 1, \dots, 6. \quad (19)$$

Now we give the solution for the preonic $\mathbf{N}=1$ supersymmetric non-Abelian flux tubes based on theory [5–7]. Dual symmetry included in our model slightly modifies theory [5–7] by consideration of the Zwanziger formalism [11, 12].

4. As it was shown in Refs. [11, 12], the aim to describe symmetrically non-dual and dual Abelian fields A_μ and $\tilde{A}_\mu$, covariantly interacting with electric $j_\mu^{(e)}$ and magnetic $j_\mu^{(m)}$ currents respectively, is realized by the following Zwanziger’s action:

$$S^{ZW} = \int d^4x \left(-\frac{1}{2} n^\mu n^\lambda \sqrt{-g} g^{\nu\rho} \right) \left(F_{\mu\nu} F_{\lambda\rho} + G_{\mu\nu} F_{\lambda\rho} + i F_{\mu\nu} \tilde{G}_{\lambda\rho} - i G_{\mu\nu} \tilde{F}_{\lambda\rho} \right), \quad (20)$$

where

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (21)$$

$$G_{\mu\nu} = \partial_\mu \tilde{A}_\nu - \partial_\nu \tilde{A}_\mu, \quad (22)$$

and

$$\tilde{F}_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F_{\rho\sigma}. \quad (23)$$

In Eq. (20) a constant unit vector n^μ denotes the direction of Dirac strings, which are frozen in time and parallel in space.

The generalized Zwanziger formalism for non-Abelian gauge theories was suggested in Refs. [12, 13] (see also references therein), and we consider the following Zwanziger-type action:

$$\begin{aligned} S^{NAZW} = & -\frac{1}{4\pi} \int \mathcal{D}\xi^\mu(s) ds \left\{ \text{Tr} \left[\left(\dot{\xi}^\mu(s) F_{\mu\nu}(x) \right) \left(\dot{\xi}^\lambda(s) F_\lambda^\nu(x) \right) \right] \dot{\xi}^{-2} \right. \\ & + \text{Tr} \left[\left(\dot{\xi}^\mu(s) G_{\mu\nu}(x) \right) \left(\dot{\xi}^\lambda(s) G_\lambda^\nu(x) \right) \right] \dot{\xi}^{-2} \\ & + \text{Tr} \left[\left(\dot{\xi}^\mu(s) F_{\mu\nu}(x) \right) \left(\dot{\xi}^\lambda(s) \tilde{G}_\lambda^\nu(x) \right) \right] \dot{\xi}^{-2} \\ & \left. + \text{Tr} \left[\left(\dot{\xi}^\mu(s) G_{\mu\nu}(x) \right) \left(\dot{\xi}^\lambda(s) \tilde{F}_\lambda^\nu(x) \right) \right] \dot{\xi}^{-2} \right\}, \quad (24) \end{aligned}$$

where $\xi^\mu(s)$ represents an arbitrary disposition of string in D -dimensional space-time ($D = 4$ in our case). The functional integral $\int \mathcal{D}\xi^\mu(s)$ introduces the sum over the string’s shape. The “tilde” denotes “dual”, but in this non-Abelian case it is not a simple “Hodge star duality” given by Eq. (23) (see [12, 13]).

In Eq. (24) we have:

$$F_{\mu\nu} = \partial_\nu A_\mu(x) - \partial_\mu A_\nu(x) + ig [A_\mu(x), A_\nu(x)], \quad (25)$$

and

$$G_{\mu\nu}(x) = \partial_\nu \tilde{A}_\mu(x) - \partial_\mu \tilde{A}_\nu(x) + i\tilde{g} [\tilde{A}_\mu(x), \tilde{A}_\nu(x)]. \quad (26)$$

Here and below we do not consider the “absorption” of the coupling constants g and $\tilde{g}$ by vector potentials A_μ and $\tilde{A}_\mu$.

5. Near the Planck scale the gauge group for preons is supersymmetric

$$[SU(6) \times U(1)] \times [\widetilde{SU(6)} \times \widetilde{U(1)}],$$

revealing the generalized dual symmetry (see [11–13] and references therein). In the non-dual sector, besides $SU(6)$ and $U(1)$ gauge bosons, preons $P = (P_1, P_2^+)^T$ (Dirac fermions and $SU(2)_R$ singlets) and other matter fields, the model contains six scalar fields $\tilde{P}$ (spreons) charged with respect to $U(1)$ gauge group and belong to the 6-plet of $SU(6)$. In general, duality leads to the 6×6 matrix (18) for spreons. Also they are $SU(2)_R$ doublets: $\tilde{P} = (\tilde{P}_1, \tilde{P}_2)$. As in Refs. [5], we introduce a zero charge scalar field f and 35×35 matrix of the complex adjoint scalar fields:

$$f = \{f^{jJ}\} \quad \text{with} \quad j, J = 1, 2, \dots, 35. \quad (27)$$

In terms of these fields the action of $\mathbf{N}=2$ supersymmetric theory can be deformed by the breaking to $\mathbf{N}=1$ terms containing mass parameters μ_1 and μ_2 for the fields f^{jJ} and f . The action takes the following form:

$$\begin{aligned} S = & S^{ZW} + S^{NAZW} + S^{(\text{matter})} \\ & + \text{other terms (topological, gauge fixing, etc.)}, \end{aligned} \quad (28)$$

where S^{ZW} and S^{NAZW} are given by Eqs. (20) and (24), and the third term in Eq. (28) is:

$$\begin{aligned} S^{(\text{matter})} = & \int d^4x \left(|D_\mu f^{jJ}|^2 + |\nabla_\mu \tilde{P}_1|^2 + |\nabla_\mu \tilde{P}_2|^2 + |\partial_\mu f|^2 + U(\tilde{P}, f^{jJ}, f) \right) \\ & + \text{other matters}, \end{aligned} \quad (29)$$

where D_μ is the covariant derivative in the adjoint representation defined as

$$D_\mu = \partial_\mu - ig_6 [A_\mu(x), \dots] - i\tilde{g}_6 [\tilde{A}_\mu(x), \dots], \quad (30)$$

while

$$\nabla_\mu = \partial_\mu - \frac{i}{\sqrt{2N}} \left(g_1 A_\mu + \tilde{g}_1 \tilde{A}_\mu \right) - i \left(g_6 A_\mu^j T^j + \tilde{g}_6 \tilde{A}_\mu^J T^J \right). \quad (31)$$

In our case $N = 6$, and T^j, T^J are $SU(6)$ generators. The charge g_1 and $(\tilde{g}_1)$ belongs to the $U(1)$ and $(\widetilde{U(1)})$ gauge group respectively and g_6 and $(\tilde{g}_6)$ is the charge of the $SU(6)$ and $(\widetilde{SU(6)})$ gauge group of theory respectively.

The potential $U(\tilde{P}, f^{jJ}, f)$ is a sum of various supersymmetric D and F terms:

$$\begin{aligned}
U(\tilde{P}, f^{jJ}, f) = & \frac{1}{2g_6^{*2}} \left(\frac{1}{g_6^{*2}} (f^{jJ})^+ f^{jJ} + \overline{\tilde{P}_1} T^j T^J \tilde{P}_1 - \tilde{P}_2 T^j T^J \overline{\tilde{P}_2} \right)^2 \\
& + \frac{g_1^{*2}}{8} \left(|\tilde{P}|^2 - N\xi \right)^2 + g_6^{*2} \left([\tilde{P}_2 T^j T^J \tilde{P}_1] + \mu_2 f^{jJ} \right)^2 \\
& + g_1^{*2} \left(\tilde{P}_2 \tilde{P}_1 + \mu_1 f \right)^2 + \frac{1}{2} \sum_{a,A}^6 \left[\left| (f + f^{jJ} T^j T^J) \tilde{P}_1^{aA} \right|^2 \right. \\
& \left. + \left| (f + f^{jJ} T^j T^J) \overline{\tilde{P}_{2,aA}} \right|^2 \right], \tag{32}
\end{aligned}$$

where we have charges:

$$g_1^{*2} = g_1^2 + \tilde{g}_1^2 \quad \text{and} \quad g_6^{*2} = g_6^2 + \tilde{g}_6^2, \tag{33}$$

which are a result of the dual symmetry.

In Eq. (32) we have a parameter ξ belonging to the Fayet-Iliopoulos D -term, which does not break $\mathbf{N}=2$ supersymmetry. The breakdown to $\mathbf{N}=1$ supersymmetric theory is realized with the help of parameters μ_1 and μ_2 (see [5]). However, the Fayet-Iliopoulos D -term triggers the spontaneous breaking of the gauge symmetry.

The action (28)–(33) gives the solution for the supersymmetric non-Abelian flux tubes in our preonic model. This solution was obtained by method developed in Ref. [5].

The vacuum expectation value (VEV) of spreons is given as

$$v = \sqrt{\xi} \gg \Lambda_4, \tag{34}$$

where Λ_4 is the 4-dimensional scale. This VEV is equal to

$$v \sim M_{Pl} \sim 10^{19} \text{ GeV}, \tag{35}$$

because we assume that spreons are condensed at the Planck scale.

Non-trivial topology (17) amounts to winding of elements of matrix (18), and we obtain string solutions:

$$\tilde{P}_{string} = v \cdot \text{diag} \left(e^{i\alpha(x)}, e^{i\alpha(x)}, \dots, 1, 1 \right), \quad \text{where } x \rightarrow \infty. \tag{36}$$

Three types of string moduli space

$$\frac{SU(6)}{SU(5) \times U(1)}, \quad \frac{SU(6)}{SU(4) \times SU(2) \times U(1)} \quad \text{and} \quad \frac{SU(6)}{SU(3) \times SU(3) \times U(1)} \tag{37}$$

give us solutions for three types of Z_6 -flux tubes which are a non-Abelian analog of ANO-strings.

6. Assuming at the ends of strings the existence of the preon P and antipreon P^c with hyper-magnetic charges $n\tilde{g}$ and $-n\tilde{g}$, respectively, we obtain six types of strings having the fluxes Φ_n quantized according to the Z_6 center group of $SU(6)$:

$$\Phi_n = n\Phi_0, \quad n = \pm 1, \pm 2, \pm 3. \quad (38)$$

Indeed, Z_6 has six group elements:

$$Z_6 = \left\{ \exp \left(2\pi \frac{n}{6} i \right) \middle| n \bmod 6 \right\}. \quad (39)$$

So far as n is given by modulo 6, the fluxes of tubes corresponding to the solutions with $n = 4, 5$ are equal to the fluxes with $n = -2, -1$, respectively (see also [6]).

String tensions of these non-Abelian flux tubes also were calculated in Refs. [5]. The minimal tension is:

$$T_0 = 2\pi\xi, \quad (40)$$

which in our preonic model is equal to:

$$T_0 = 2\pi v^2 \sim 10^{38} \text{ GeV}^2. \quad (41)$$

Such an enormously large tension means that preonic strings have almost infinitely small $\alpha' \rightarrow 0$, where $\alpha' = 1/(2\pi T_0)$ is a slope of trajectories in string theory [1]. Three types of the preonic k -strings have the following tensions:

$$T_k = kT_0, \quad \text{where } k = 1, 2, 3. \quad (42)$$

If the Fayet-Iliopoulos term ξ vanishes in the supersymmetric theory of preons, then spreon condensate vanishes too, and the theory is in the Coulomb phase. But for a non-vanishing ξ the spreons develop their VEV, and the theory is in the Higgs phase. Then hyper-magnetic charges of preons and antipreons are confined by six strings which are oriented in opposite directions. By this reason, six strings have only three different tensions (42).

Also preonic strings are enormously thin. Indeed, the thickness of the flux tube depends on the mass of the dual gauge boson $\tilde{A}_\mu$ acquired in the confinement phase:

$$m_V = gv. \quad (43)$$

As it is shown below, in the region of energies AB near the Planck scale, where spreons are condensed (see Fig. 2) we have:

$$\alpha = \frac{g^2}{4\pi} \approx 1, \quad g \approx 2\sqrt{\pi} \approx 3.5, \quad (44)$$

and the thickness of preonic strings given by the radius R_{str} of the flux tubes is very small:

$$R_{str} \sim \frac{1}{m_V} \sim \frac{1}{gv} \sim 10^{-18} \text{ GeV}^{-1}. \quad (45)$$

Such infinitely narrow non-Abelian supersymmetric flux tubes remind us superstrings of Superstring theory. Having in our preonic model supersymmetric strings with $\alpha' \rightarrow 0$ we obtain, according to the description [1], only massless ground states: spin 1/2 fermions (quarks and leptons), spin 1 hyper-gluons and spin 2 massless graviton, as well as their superpartners. The excited states belonging to these strings are not realized in our world as very massive: they have mass $M > M_{Pl}$.

7. The hyper-flavor “horizontal” symmetry was suggested first in Refs. [14]. The previous Section gives a demonstration of a very specific type of the “horizontal symmetry”: three, and only three, generations of fermions and bosons present in the superstring-inspired flipped E_6 theory, and also in each step given by Fig. 1 up to the Standard Model. This number “3” is explained by the existence of three values of hyper-magnetic fluxes, which bind hyper-magnetic charges of preons-dyons. At the ends of these preonic strings there are placed hyper-magnetic charges $\pm\tilde{g}_0$, or $\pm 2\tilde{g}_0$, or $\pm 3\tilde{g}_0$, where $\tilde{g}_0$ is the minimal hyper-magnetic charge. Then all bound states of Fig. 3 form three generations – three 27-plets of E_6 corresponding to the three different tube flux values. We also obtain the three types of gauge bosons A_μ^i ($i = 1, 2, 3$ is the index of generations) belonging to the $27 \times \overline{27} = 1 + 78 + 650$ representations of E_6 . Fig. 4 illustrates the formation of such hyper-gluons: Fig. 4(a) corresponds to the composite singlet, Fig. 4(b) and Fig. 4(c) correspond to the adjoint 78-plet and 650-plet of hyper-gluons, respectively.

Such a description predicts the Family replicated gauge group of symmetry $[E_6]^3$ for quark-leptons and bosons, which works near the Planck scale. Here the number of families is equal to the number of generations $N_g = 3$. We assume that $[E_6]^3$ for quark-leptons takes place in the region of energies $M_{FR} \leq \mu \leq M_{crit}$, where the scales M_{FR} and M_{crit} are indicated in Fig. 2 by points D and A , respectively.

The breakdown $[E_6]^3 \rightarrow E_6$ at the scale M_{FR} is provided by several Higgses. The analogous mechanism is described in reviews [15] and references therein. Indeed, in the Family replicated gauge group we have three types of gauge bosons A_μ^i , which produce linear combinations:

$$A_{\mu,diag}^{(i)} = C_1^{(i)} A_\mu^{(1st\ fam.)} + C_2^{(i)} A_\mu^{(2nd\ fam.)} + C_3^{(i)} A_\mu^{(3rd\ fam.)}, \quad i = 1, 2, 3. \quad (46)$$

The combination:

$$A_{\mu,diag} = \frac{1}{\sqrt{3}} (A_\mu^{(1st\ fam.)} + A_\mu^{(2nd\ fam.)} + A_\mu^{(3rd\ fam.)}) \quad (47)$$

is massless one, which corresponds to the 78 adjoint representation of hyper-gluons of E_6 . Two other combinations are massive and exist only in the region of energies $\mu \geq M_{FR}$.

The assumption that only $[E_6]^3$ for quark-leptons exists in Nature may be not valid: we are not sure that the Family replicated gauge groups $[SO(10)]^3$, $[SU(5)]^3$, $[SMG \times U(1)_{(B-L)}]^3$, or $[SMG]^3$ do not survive at lower energies $\mu < M_{FR}$. Here we have used the following notation: SMG is the Standard Model group $SU(3)_C \times SU(2)_L \times U(1)_Y$.

The values $\alpha^{-1}(M_{crit})$ and $\alpha^{-1}(\widetilde{M}_{crit})$ can be calculated by the method developed in Ref. [16]. If $SU(N)$ group is broken by Abelian scalar particles belonging to its Cartan $U(1)^{N-1}$ subalgebra, then $SU(N)$ critical coupling constant α_N^{crit} is given by the following expression in the one-loop approximation (see [13, 16]):

$$\alpha_N^{crit} \approx \frac{N}{2} \sqrt{\frac{N+1}{N-1}} \alpha_{U(1)}^{crit}, \quad (48)$$

where $\alpha_{U(1)}^{crit}$ is the critical coupling constant for the Abelian $U(1)$ theory.

Here it is necessary to distinguish E_6 gauge group of symmetry for preons from E_6 for quark-leptons. The points A and B of Fig. 2 respectively correspond to the breakdowns of $[E_6]^3$ and $[\widetilde{E}_6]^3$ to the region AB of spreon condensation. There we have the breakdown (see Ref. [3]) of the preon (one family) E_6 (or $\widetilde{E}_6$):

$$E_6 \rightarrow SU(6) \times U(1) \quad \left(\text{or} \quad \widetilde{E}_6 \rightarrow \widetilde{SU(6)} \times \widetilde{U(1)} \right),$$

and the scale M_{crit} (or $\widetilde{M}_{crit}$) is the scale of breaking.

In our preonic model the group $SU(6)$ is broken by condensed Abelian scalar spreons-dyons belonging to the Cartan subalgebra $U(1)^5$. Then for the one family of preons we have:

$$\alpha_6^{crit} \approx 3.55 \alpha_{U(1)}^{crit}, \quad (49)$$

according to Eq. (48) for $N = 6$.

The behaviour of the effective fine structure constants in the vicinity of the phase transition point ‘‘Coulomb-confinement’’ was investigated in the compact lattice $U(1)$ theory by Monte Carlo method [18]. The following result was obtained:

$$\alpha_{lat.U(1)}^{crit} \approx 0.20 \pm 0.015, \quad \tilde{\alpha}_{lat.U(1)}^{crit} \approx 1.25 \pm 0.010. \quad (50)$$

The calculation of the critical coupling constants in the Higgs scalar monopole (dyon) model of dual $U(1)$ theory [16, 17] gave the following result:

$$\alpha_{U(1)}^{crit} \approx 0.21, \quad \tilde{\alpha}_{U(1)}^{crit} \approx 1.20 \quad - \quad \text{in the Higgs monopole model}, \quad (51)$$

and

$$\alpha_{U(1)}^{crit} \approx 0.19, \quad \tilde{\alpha}_{U(1)}^{crit} \approx 1.29 \quad - \quad \text{in the Higgs dyon model}. \quad (52)$$

According to Eqs. (50)–(52), the condensation of spreons leads to the following critical constants:

$$\alpha_6^{crit} \approx 3.55 \cdot 0.2 \approx 0.71, \quad \tilde{\alpha}_6^{crit} = (\alpha_6^{crit})^{-1} \approx 1.41. \quad (53)$$

For the point C shown in Fig. 2 we have:

$$\alpha_6^{(one\ fam.)}(M_{Pl}) = 1. \quad (54)$$

This result for the one family of preons confirms the estimate (44). Here we have used the Dirac relation for non-Abelian theories (see explanation in Ref. [13]):

$$g\tilde{g} = 4\pi n, \quad n \in Z, \quad \alpha\tilde{\alpha} = 1. \quad (55)$$

From the phase transition result (53) for the one family of preons, we obtain the following inversed critical coupling constant for $[E_6]^3$ at the phase transition point A (see reviews [15]):

$$\alpha_A^{-1}(M_{crit}) = 3\tilde{\alpha}_6^{crit}(M_{crit}) \approx 3 \cdot 1.41 \approx 4.23. \quad (56)$$

At the phase transition point B , when we have the breakdown of $[\tilde{E}_6]^3$ into the confinement phase, the one-family value $\tilde{\alpha}_6^{crit}(\tilde{M}_{crit}) \approx 0.71$ gives the following result:

$$\alpha_B^{-1}(\tilde{M}_{crit}) \approx 3 \cdot 0.71 \approx 2.13. \quad (57)$$

The values (56) and (57) were used for the construction of the curve AB presented in Fig. 2. The point C corresponds to the Planck scale and $\alpha^{-1}(M_{Pl}) = 3$, according to Eq. (54).

The condensation of spreons at the Planck scale predicts the existence of the second minimum of the effective potential $V_{eff}(\mu)$ at the scale $\mu = M_{Pl}$. The behaviour of this potential and its relation with the Multiple Point Principle – theory of degenerate vacua by D.L. Bennett, C.D. Froggatt and H.B. Nielsen (see reviews [15] and references therein) will be considered in our future investigation.

Fig. 2 shows two points A and B near the Planck scale. The point A corresponds to the breakdown of E_6 for preons according to the chain (16). The group E_6 is broken in the region of energies $\mu \geq M_{crit}$ producing hyper-electric strings between preons. The point B in Fig. 2 indicates the scale $\tilde{M}_{crit}$ corresponding to the breakdown $\tilde{E}_6 \rightarrow \widetilde{SU(6)} \times \widetilde{U(1)}$. At the point B hyper-magnetic strings are produced and exist in the region of energies $\mu \leq \tilde{M}_{crit}$ confining hyper-magnetic charges of preons. As a result, in the region $\mu \leq \tilde{M}_{crit}$ we see quark-leptons with charges ng ($n \in Z$), but in the region $\mu \geq \tilde{M}_{crit}$ monopolic “quark-leptons” – particles with dual charges $m\tilde{g}$ ($m \in Z$) – may exist. Since $\tilde{M}_{crit} > M_{Pl}$, monopoles are absent in our world.

In the region of energies $M_{crit} \leq \mu \leq \tilde{M}_{crit}$ around the Planck scale, both hyper-electric and hyper-magnetic strings come to play: preons, quarks and monopolic “quarks” are totally confined giving heavy neutral particles with mass $M \sim M_{Pl}$, but closed strings – gravitons – survive there.

The dotted curve in Fig. 2 describes the running of $\alpha^{-1}(\mu)$ for monopolic “quark-leptons” created by preons, which are bound by supersymmetric hyper-electric non-Abelian flux tubes. We assume that such monopoles do not exist in our world, however, they can play an essential role in the Universe vacuum (Cosmological Constant) [19].

Conclusions

- i. In the present paper starting with an idea that the most realistic model based on the superstring theory is the ‘flipped’ E_6 gauge group of symmetry, we have assumed that at high energies $\mu > 10^{16}$ GeV there exists the following chain of the flipped models:

$$SU(3)_C \times SU(2)_L \times U(1)_Y \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X \rightarrow$$

$$SU(5) \times U(1)_X \rightarrow SU(5) \times U(1)_{Z1} \times U(1)_{X1} \rightarrow SO(10) \times U(1)_{X1} \rightarrow E_6,$$

with the flipped E_6 final unification. We have chosen such Higgs boson contents of the $SU(5)$ and $SO(10)$ gauge groups, which give the flipped E_6 at the scale $\sim 10^{18}$ GeV and the decreased running of $\alpha^{-1}(\mu)$ near the Planck scale.

- ii. We have shown that the final unification E_6 assumes the existence of three 27-plets of the flipped E_6 gauge group of symmetry.
- iii. Suggesting $\mathbf{N} = 1$ supersymmetric $E_6 \times \widetilde{E}_6$ preonic model of composite quark-leptons and bosons, we have assumed that preons are dyons confined by hyper-magnetic strings in the region of energies $\mu \lesssim M_{Pl}$. This approach is an extension of the old idea by J. Pati to use the strong magnetic forces which may bind preons-dyons in composite particles – quark-leptons and bosons. Our model is based on the recent theory of composite non-Abelian flux tubes in SQCD [5–7].
- iv. Considering the breakdown of E_6 (or $\widetilde{E}_6$) at the Planck scale into the $SU(6) \times U(1)$ (or $\widetilde{SU}(6) \times \widetilde{U}(1)$) gauge group we have shown that the six types of k -strings – $\mathbf{N} = 1$ supersymmetric non-Abelian flux tubes – are created by the condensation of spreons near the Planck scale.
- v. It was shown that the six types of strings-tubes having six fluxes quantized according to the Z_6 center group of $SU(6)$:

$$\Phi_n = n\Phi_0, \quad n = \pm 1, \pm 2, \pm 3,$$

create three types of k -strings with tensions:

$$T_k = kT_0, \quad k = 1, 2, 3,$$

which produce three (and only three) generations of composite quark-leptons and bosons. We have obtained a specific type of the “horizontal symmetry” explaining a flavor.

- vi. It was investigated that in the present model preonic strings are very thin, with radius

$$R_{str} \sim 10^{-18} \text{ GeV}^{-1},$$

and their tension is enormously large:

$$T_0 \sim 10^{38} \text{ GeV}^2.$$

- vii. The model predicts the existence of three families of 27-plets and also gauge bosons A_μ^i (with $i = 1, 2, 3$) belonging to the 78-plets of E_6 . Then near the Planck scale we have the Family replicated gauge group of symmetry $[E_6]^3$. In the present paper we have assumed that the breakdown $[E_6]^3 \rightarrow E_6$ occurs near the Planck scale leading to the E_6 unification at the scale $\sim 10^{18}$ GeV.
- viii. We have considered that the condensation of spreons-dyons near the Planck scale gives the phase transitions at the scales M_{crit} and $\widetilde{M}_{crit}$ shown in Fig. 2. These scales (points A, B of Fig. 2) correspond to the breakdown of E_6 and $\widetilde{E}_6$ for preons, respectively:

$$E_6 \rightarrow SU(6) \times U(1) \quad \text{and} \quad \widetilde{E}_6 \rightarrow \widetilde{SU(6)} \times \widetilde{U(1)}.$$

- ix. It was investigated that hyper-magnetic strings are produced and exist at $\mu \leq \widetilde{M}_{crit}$, and hyper-electric strings are created and exist at $\mu \geq M_{crit}$. As a result, in our world we have quark-leptons and gauge bosons A_μ in the region of energies $\mu \lesssim M_{Pl}$, but monopolic “quark-leptons” and dual gauge fields $\widetilde{A}_\mu$ exist in the region $\mu \gtrsim M_{Pl}$ (at the trans-Planckian scales, see [19]). We have calculated the critical values of gauge coupling constants at the scales M_{crit} and $\widetilde{M}_{crit}$:

$$\alpha^{-1}(M_{crit}) \approx 4.23, \quad \text{and} \quad \alpha^{-1}(\widetilde{M}_{crit}) \approx 2.13.$$

Acknowledgements: C.R.D. thanks a lot Prof. J. Pati for interesting discussions during the Conference WHEPP-9 (Bhubaneswar, India, January, 2006). L.L. sincerely thanks the Institute of Mathematical Sciences (Chennai, India) and personally Director of IMSc Prof. R. Balasubramanian and Prof. N.D. Hari Dass for the wonderful hospitality and financial support. The authors also deeply thank Prof. J.L. Chkareuli, Prof. F.R. Klinkhamer and Prof. G. Rajasekaran for fruitful discussions and advices.

This work was supported by the Russian Foundation for Basic Research (RFBR), project N° 05-02-17642.

References

- [1] J. Schwarz, Phys.Rept. **89**, 223 (1982); M.B. Green, Surv.High.En.Phys. **3**, 127 (1984); M.B. Green and J.H. Schwarz, Phys.Lett. **B149**, 117 (1984); *ibid.*, **B151**, 21 (1985); D.J. Gross, J.A. Harvey, E. Martinec and R. Rohm, Phys.Rev.Lett. **54**, 502 (1985); Nucl.Phys. **B256**, 253 (1985); *ibid.*, **B267**, 75 (1986).
- [2] J.C. Pati, Phys.Lett. **B98**, 40 (1981); J.C. Pati, A. Salam and J. Strathdee, Nucl.Phys. **B185**, 416 (1981).
- [3] C.R. Das and L.V. Laperashvili, Phys.Rev. **D74**, 035007 (2006); arXiv: hep-ph/0605161.
- [4] C.R. Das and L.V. Laperashvili, *Composite model of quark-leptons and duality*, a talk prepared for the 17th DAE-BRNS High Energy Physics Symposium, IIT, Kharagpur, India, 11-15 December, 2006; arXiv: hep-ph/0606042.

- [5] M. Shifman and A. Yung, Phys.Rev. **D70**, 025013 (2004), arXiv: hep-th/0312257; ibid., **D70**, 045004 (2004), arXiv: hep-th/0403149; ibid., **D72**, 085017 (2005), arXiv: hep-th/0501211; A. Gorsky, M. Shifman and A. Yung, Phys.Rev. **D71**, 045010 (2005), arXiv: hep-th/0412082; V. Markov, A. Marshakov and A. Yung, Nucl.Phys. **B709**, 267 (2005), arXiv: hep-th/0408235; R. Auzzi, M. Shifman and A. Yung, Phys.Rev. **D73**, 105012 (2006), arXiv: hep-th/0511150.
- [6] M.A.C. Kneipp, Phys.Rev. **D69**, 045007 (2004); arXiv: hep-th/0308086.
- [7] Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, Phys.Rev. **D71**, 065018 (2005), arXiv: hep-th/0405129; M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, Phys.Rev.Lett. **96**, 161601 (2006), arXiv: hep-th/0511088; M. Eto, T. Fujimori, Y. Isozumi, M. Nitta, K. Ohashi, K. Ohta and N. Sakai, Phys.Rev. **D73**, 085008 (2006), arXiv: hep-th/0601181; M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, Phys.Rev. **D73**, 125008 (2006), arXiv: hep-th/0602289.
- [8] C.R. Das and L.V. Laperashvili, *Seesaw scales and the steps from the Standard Model towards superstring-inspired flipped E_6* ; arXiv: hep-ph/0604052.
- [9] C.R. Das, C.D. Froggatt, L.V. Laperashvili and H.B. Nielsen, Mod.Phys.Lett. **A21**, 1151 (2006); arXiv: hep-ph/0507182.
- [10] M. Chaichian, J.L. Chkareuli and A. Kobakhidze, Phys.Rev. **D66**, 095013 (2002); arXiv: hep-ph/0108131.
- [11] D. Zwanziger, Phys.Rev. **176**, 1489 (1968); ibid., Phys.Rev. **D3**, 880 (1971); R.A. Brandt, F. Neri and D. Zwanziger, Phys.Rev. **D19**, 1153 (1979); F.V. Gubarev, M.I. Polikarpov and V.I. Zakharov, Phys.Lett. **B438**, 147 (1998), arXiv: hep-th/9805175; L.V. Laperashvili and H.B. Nielsen, Mod.Phys.Lett. **A14**, 2797 (1999), arXiv: hep-th/9910101.
- [12] L.V. Laperashvili, *Generalized dual symmetry of non-Abelian theories*, arXiv: hep-th/0211227; A. Noguchi and A. Sugamoto, *Dynamical origin of duality between gauge theory and gravity*, in: TSPU Vestnik **44N7**, 2004, p.59., arXiv: hep-th/0408045.
- [13] C.R. Das, L.V. Laperashvili and H.B. Nielsen, *Generalized dual symmetry of non-Abelian theories and the freezing of α_s* , to appear in Int.J.Mod.Phys. **A**, (2006); arXiv: hep-ph/0511267.
- [14] J.L. Chkareuli, JETP Lett. **32**, 671 (1980) [Pisma Zh.Eksp.Teor.Fiz. **32**, 684 (1980)]; Z.G. Berezhiani and J.L. Chkareuli, Sov.J.Nucl.Phys. **37**, 618 (1983) [Yad. Fiz. **37**, 1043 (1983)].
- [15] C.D. Froggatt, L.V. Laperashvili, H.B. Nielsen and Y. Takanishi, *Family Replicated Gauge Group Models*, in: Proceedings of the “Fifth International Conference Symmetry in Nonlinear Mathematical Physics”, Kiev, Ukraine, 23-29 June, 2003, Ed. by

- A.G. Nikitin, V.M. Boyko, R.O. Popovich and I.A. Yehorchenko (Institute of Mathematics of NAS of Ukraine, Kiev, 2004), Vol.50, Part 2, p.737, arXiv: hep-ph/0309129; L.V. Laperashvili, Phys.At.Nucl. **57**, 471 (1994) [Yad.Fiz. **57**, 501 (1994)]; ibid., **59**, 162 (1996) [Yad.Fiz. **59**, 172 (1996)].
- [16] L.V. Laperashvili, H.B. Nielsen and D.A. Ryzhikh, Int.J.Mod.Phys. **A16**, 3989 (2001), arXiv: hep-th/0105275; L.V. Laperashvili and H.B. Nielsen, Int.J.Mod.Phys. **A16**, 2365 (2001), arXiv: hep-th/0010260.
- [17] C.R. Das and L.V. Laperashvili, *Phase transition in the Higgs model of scalar dyons*, to appear in Mod.Phys.Lett. **A** (2006); arXiv: hep-ph/0511067.
- [18] J. Jersak, T. Neuhaus and P.M. Zerwas, Phys.Lett. **B133**, 103 (1983); Nucl.Phys. **B251**, 299 (1985); J. Jersak, T. Neuhaus and H. Pfeiffer, Phys.Rev. **D60**, 054502 (1999), arXiv: hep-lat/9903034.
- [19] C. Froggatt, L. Laperashvili, R. Nevzorov and H.B. Nielsen, Phys.Atom.Nucl.**67**, 582 (2004) [Yad.Fiz. **67**, 601 (2004)], arXiv: hep-ph/0310127; F.R. Klinkhamer and G.E. Volovik, JETP Lett. **81**, 551 (2005), [Pisma Zh.Eksp.Teor.Fiz. **81**, 683 (2005)] arXiv: hep-ph/0505033.




Fig. 1: This figure presents the running of the inversed gauge coupling constants $\alpha_i^{-1}(x)$ for $i = 1, 2, 3, X, Z, X1, Z1, 5, 10$ of the chain $SU(3)_C \times SU(2)_L \times U(1)_Y \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X \rightarrow SU(5) \times U(1)_X \rightarrow SU(5) \times U(1)_{Z1} \times U(1)_{X1} \rightarrow SO(10) \times U(1)_{X1} \rightarrow E_6$ corresponding to the breakdown of the flipped $SU(5)$ to the supersymmetric (MSSM) $SU(3)_C \times SU(2)_L \times U(1)_Z \times U(1)_X$ gauge group of symmetry with Higgs bosons belonging to the $5_h + \bar{5}_h, 10_H + \bar{10}_H$, 24-dimensional adjoint A and higher representations of the flipped $SU(5)$. The final unification is E_6 at the supersuperGUT scale $M_{SSG} \approx 10^{18}$ GeV with $\alpha^{-1}(M_{SSG}) \approx 22$ for $M_{SUSY} = 10$ TeV and seesaw scale $M_{SS} = 10^{11}$ GeV.




Fig. 2: The figure illustrates a qualitative description of the running of $\alpha^{-1}(x)$ near the Planck scale predicted by the present preonic model. The region AD corresponds to the Family replicated gauge group of symmetry $[E_6]^3$, which comes at the scale M_{FR} given by point D . The point A at the scale of energy $\mu = M_{crit}$ shows that hyper-electric preonic strings exist for $\mu \geq M_{crit}$. The point B corresponds to the scale $\mu = \tilde{M}_{crit}$ and indicates that hyper-magnetic preonic strings exist for $\mu \leq \tilde{M}_{crit}$. The curve AB corresponds to the region of energies, where spreons are condensed near the Planck scale giving both, hyper-electric and hyper-magnetic, preonic strings. For $\mu \geq \tilde{M}_{crit}$ we have the running of $\alpha^{-1}(x)$ for monopolic “quark-leptons”. The point C corresponds to the Planck scale and gives $\alpha^{-1}(M_{Pl}) = 3$.




Fig. 3: Preons are bound by hyper-magnetic strings: (a,b) correspond to the string configurations of composite particles belonging to the 27-plet of E_6 gauge group of symmetry; (c) represents a closed string describing a graviton.




Fig. 4: Vector gauge bosons belonging to the $1 + 78 + 650$ representations of E_6 . Gluons are composite objects created by fermionic preons P, P^c : (a) corresponds to the singlet; (b) is the 78-plet and (c) is the 650-plet of E_6 .